

B. B. BROWNE (B.B.)

A PAPER

ON

SUBINVOLUTION OF THE UTERUS.

BY

B. B. BROWNE, M. D.,

307 Madison Avenue
BALTIMORE, MD.

[Read before the Clinical Society of Baltimore, March 2, 1877.]


FROM THE JUNE NO. RICHMOND AND LOUISVILLE MEDICAL JOURNAL

LOUISVILLE, KY.:

RICHMOND AND LOUISVILLE MEDICAL JOURNAL BOOK AND JOB STEAM PRINT,

104 Green Street, 2d door west of Post-office.

1877.



A PAPER

ON

SUBINVOLUTION OF THE UTERUS.

BY

B. B. BROWNE, M. D.,

BALTIMORE, MD.

[Read before the Clinical Society of Baltimore, March 2, 1877.]

FROM THE JUNE NO. RICHMOND AND LOUISVILLE MEDICAL JOURNAL.

LOUISVILLE, KY.:

RICHMOND AND LOUISVILLE MEDICAL JOURNAL BOOK AND JOB STEAM PRINT,

104 Green Street, 2d door west of Post-office.

1877.

SUBINVOLUTION OF THE UTERUS.

Normal puerperal convalescence includes the restoration of the pelvic organs to their normal state, which during gestation and parturition have been the seat of extraordinary modification in tissue, function and position. During the period of uterogestation the uterus enlarges from about three inches in length and one and three-quarters in breadth to twelve or fifteen in length and nine or ten in breadth. It increases from about two ounces in weight to twenty-five or thirty ounces. Its cavity before impregnation is less than one cubic inch, while at the full term of pregnancy it is extended to above four hundred cubic inches, and the surface of the organ increases from about five or six square inches to nearly three hundred and fifty square inches. The reduction of the uterus after delivery to its normal size, its involution as it is termed, takes place by muscular contractions, fatty transformation of its component fibres and absorption. The celerity of its involution in the puerperal state is more striking and remarkable than the celerity of its evolution during the pregnant state. For while the uterus takes forty weeks to acquire its dimensions at the full term pregnancy, it requires only from six to eight weeks to decrease to the small size of the organ in its unimpregnated condition.

If the conditions necessary for involution are interfered with or impeded subinvolution will be the result. This condition may be caused by a great number of diseases and accidents that are liable to occur during the puerperal state. For, however easy and healthy the pregnancy and delivery may have been, however happily they may have run their course, yet every lying-in woman is in a high degree predisposed to puerperal affections. The pathetic utterances of an old gynæcologist, Chambon de Montaux, that woman is a being whom nature has, through all ages, allowed to walk upon the edge of an abyss that is ever ready to swallow her up, surely has a peculiar sig-

nificance in childbed. For who is not familiar with those extremely rapid diseases, which within a few days or even hours, carry off the strongest women in their very prime, so that with amazement and terror he beholds their sudden end—the instantaneous and unexpected disappearance from the family circle; and of that more numerous class who escape this fatal termination, how great is the number of those who lead a life of living death, the result of puerperal diseases?

The puerperal diseases causing subinvolution of the uterus may be divided into two classes: 1. Those occurring during parturition. 2. Those occurring during the period of involution or the six or eight weeks following labor. Among the first class may be enumerated: Laceration of the cervix uteri and contusions of the uterine tissue, often caused by the maladministration of ergot to bring on uterine contractions before the os is sufficiently dilated and when the waters have already escaped; portions of retained placenta and coägula interfering with the proper contraction of the placental site and uterus; partial or complete laceration of the perineum and vestibular lacerations. Among the second class we may enumerate secondary hæmorrhages, prolonged retention of urine, constipation, improper application of the binder, rising too soon after confinement, displacement of the uterus, puerperal fever, and the non-performance of the function of lactation.

The natural function of lactation is the best and most healthy stimulus to uterine involution. The application of the infant to the breast causes contraction of the uterus. Lactation is the physiological compliment to parturition, and by causing a derivation of activity to a distant organ, tends to promote rest in the pelvis. Dr. Barnes considers the increasing neglect of the function of lactation to be a prolific cause of uterine disease. He says that this neglect does not always arise from indifference to maternal duties, or the fancied more imperative duties of social life, but that the inability to suckle is in numerous cases real. The system, the breasts want the power, the capacity to secrete milk, showing the unfitness of women nursed in luxury to carry out in its completeness the function of reproduction.

Laceration of the cervix uteri from parturition causes subin-

volution of the uterus in lateral laceration, in consequence of the irritation of the flaps which separate when the woman assumes the upright position. Dr. Emmet states that if this accident occurs from very rapid labor or from the use of forceps with considerable traction, the perineum is at the same time frequently ruptured, and with this there is a want of proper support to the uterus; it gets down upon the floor of the pelvis, and with the prolapsus there is also some degree of retroversion, so that when the woman attempts to walk the flaps are forced far apart, the posterior lip catching in the cul-de-sac and the anterior lip is crowded forward.

In subinvolution from this cause there will be more profuse leucorrhœa, menorrhagia, backache, pains down the limbs, difficulty in walking, and the uterus will be more congested and irritable.

Displacements of the uterus are a frequent cause of impeded involution, for when either retroversion, retroflexion or prolapsus occur, free circulation through the uterine vessels is necessarily interrupted, a sufficient amount of arterial blood may pass into the organ, but the return by the veins is obstructed by the tortuous course and angulation produced by the displacement.

The evil effects of an incomplete abortion in causing subinvolution are either immediate or more remote. The immediate risk is in the occurrence of hæmorrhage and the retention of fragments of the placenta and clots in the uterus; when a foreign body is thus retained, the uterus becomes decidedly hypertrophied, so that when it is finally evacuated the walls may remain permanently thickened and the cavity enlarged. When this condition exists the organ frequently becomes displaced. Such imperfect involution may easily be the starting point of other morbid changes, and many women who come under treatment for uterine diseases can trace back the commencement of their suffering to an abortion in the early months of gestation. Another cause of subinvolution of the uterus is quickly repeated pregnancy, where before the uterus is thoroughly renovated it is called upon again to undergo development.

Retarded in its metamorphosis by any such influences, the uterus does not undergo perfect involution, the fatty degenera-

tion and absorption of its muscular fibres are incompletely performed, or newly-formed tissues—muscular elements, and particularly connective tissue—are developed to such a degree as to leave the uterus considerably larger than normal. In many cases of areolar hyperplasia from subinvolution the uterus is bound down by adhesions to the pelvic viscera.

In order to avoid a more tedious detail of the various causes and effects, complications, symptoms and treatment of subinvolution of the uterus, I will give the record of the following cases:

CASE I.—June, 1875. Mrs. H.; aged thirty-three; married ten years; has had one child eight years ago; no miscarriage since; has a great deal of pain in the back and a feeling of discomfort in the lower part of the abdomen, with a sensation of weight and bearing down of the uterus, considerable pain in the lower part of the bowel and irritability of the bladder, and pain and aching extending down the limbs; a very profuse leucorrhœa. During menstruation the pain and disagreeable feelings are increased in severity and are accompanied by intense headaches lasting two or three days; the trouble has continued since the birth of the child. The uterus was enlarged, retroverted and hypertrophied, caused by subinvolution and areolar hyperplasia; there was also enlargement of the cervix with ulceration and erosion. As soon as the ulceration and congestion were somewhat reduced a suitable pessary was adjusted to support the organ in its normal position. After about three weeks' appropriate treatment the uterus was reduced to its normal size. She was relieved of her suffering and discharged cured.

CASE II.—June 25, 1875. Mrs. G.; aged thirty; married nine years; four children and one miscarriage; last child five months old. Suffering from general debility, pain and weakness in the back, heaviness and weighty feeling in the lower part of the abdomen, irritability of the bladder and leucorrhœa. Attributes her trouble to her first confinement, when she got up too soon and took cold, and had a chill and fever afterwards. Uterus large and flabby, somewhat prolapsed and retroverted—relieved by treatment, afterwards became pregnant, and expects to be confined about the middle of March.

CASE III.—April, 1876. Mrs. E.; aged twenty-eight; married six years; two children, youngest six months old; has been suffering from uterine hæmorrhages since the birth of her last child. Pressing-down pains in the lower part of her abdomen, difficult micturition and excessive weakness. Uterus large and flabby, and bleeds upon the slightest touch, is prolapsed, measures four and a half inches with the probe. Diagnosis retroversion with hypertrophy and areolar hyperplasia from subinvolution. The hæmorrhages were checked by the administration of a combination of ergot, digitalis and tincture of iron, with occasional doses of bromide of potash. By this treatment the size of the organ was also diminished. For a time it was necessary for her to wear a pessary to prevent the pressure of the cervix uteri upon the neck of the bladder, after which she was discharged cured.

CASE IV.—February 27, 1877. Mrs. C.; aged thirty-six; widow; married eighteen years ago and had one child, which was delivered with instruments after a protracted labor; her convalescence was slow, and she has suffered ever since with severe pains in her back and left side, with dysmenorrhœa and leucorrhœa; has had five or six miscarriages since the birth of her child; they have always occurred at the third month. The uterus was retroflexed with enlargement of the fundus, which was bound down by adhesions on the left side. No doubt this condition was the result of subinvolution occurring after the first pregnancy, and the adhesions by preventing the uterus from rising out of the pelvis were the cause of her numerous abortions.

PERIODICAL MEDICAL LITERATURE.

Interesting to Physicians.

Every physician who is just to his patients and to himself must read faithfully and regularly the periodical medical literature of the day. It is only thus that he can learn quickly and correctly all of the improvements constantly taking place in every department of his Profession. To all who thus seek to make themselves most efficient in practice and most successful in competition, the undersigned offers the

Richmond and Louisville Medical Journal.

This work is issued monthly, and has now reached its twenty-third volume. It is clearly printed on rose-tinted paper, which is not only agreeable to the eye, but safe, as the superfluous rays of light are absorbed by it, and are not painfully and injuriously reflected. The pages of the Journal have been increased in size, and there are now issued every year two volumes, making an aggregate of one thousand one hundred and fifty-two (1,152) pages. The cost of both volumes is but five dollars (\$5.00) annually, with twenty-five cents added for the yearly postage. Medical works of this size cost usually double the price mentioned, and are issued on paper inferior to that on which the Journal is printed. This Journal has always embodied the best articles from the best physicians in this country. Its selections are made from original European journals, and from the best medical periodicals of this country. Its exchange list is large, and includes not only all American journals, but the best of those published abroad. The reader obtains thus, every month, a condensed report of all that is valuable in American and European medicine. In addition, there is furnished the professional proceedings of medical societies; reports of interesting clinical cases; letters from European and American medical centres; reviews of all the new works; miscellaneous extracts; the medical news of the day; chemical and pharmaceutical literature; and editorial comments upon topics of current interest. The circulation of this Journal is much larger than that of any medical journal in the South, and the work is read by the leading members of the Profession. Specimen numbers 25 cents each.

There is also offered to the medical public, by the undersigned, the

AMERICAN MEDICAL BI-WEEKLY.

This Journal (as the American Medical Weekly) was founded July 4th, 1874, and is now permanently established. It is mailed invariably every other Friday night (there has not been a single failure), and reaches its subscribers regularly. It is printed on paper similar to that used for the Monthly. Its exchange list is equally large. Its pages are made up in double columns, and the subscriber receives this enlarged work for the small sum of three dollars (\$3.00) annually, with an addition of ten cents for the year's postage. The Bi-Weekly is issued in two volumes yearly; each volume having a complete index. Medical works of such size cost fully twice the price asked for this Journal. It contains original articles; extracts from foreign and domestic journals; reviews of new books, pamphlets, monographs, etc.; correspondence from European and American medical centres; the news of the day fresh and comprehensive; and editorial matter. Under no circumstances is the same material used for the two Journals; they are entirely different in the matter presented.

These Journals are owned and edited solely by the undersigned, and are therefore managed with consistency and care. They are independent of all corporations, and are strictly organs of the Medical Profession of this country.

Contributions for both Journals are respectfully requested. Those who have not the time or inclination to write formal articles can furnish matter of value in the form of letters. Every one observes something of interest, and all such facts should be given to the Profession.

E. S. GAILLARD, M. D.,

168 Second Street, Louisville, Ky.

